

# Using connectivity metrics and niche modelling to explore the occurrence of the northern crested newt *Triturus cristatus* (Amphibia, Caudata) in a traditionally managed landscape

TIBOR HARTEL<sup>1,2,3\*</sup>, SZILÁRD NEMES<sup>1</sup>, KINGA ÖLLERER<sup>4</sup>, DAN COGĂLNICEANU<sup>5</sup>, COSMIN MOGA<sup>1</sup> AND JAN W. ARNTZEN<sup>3</sup>

<sup>1</sup>Mihai Eminescu Trust, Cojocariilor 10, 545400 Sighisoara, Romania, <sup>2</sup>Babeş-Bolyai University, Department of Taxonomy and Ecology, Clinicilor 5–7, 400006 Cluj-Napoca, Romania, <sup>3</sup>Research Department Terrestrial Zoology, Netherlands Centre for Biodiversity Naturalis, PO Box 9517, 2300 RA Leiden, The Netherlands, <sup>4</sup>Institute of Biology, Romanian Academy, Splaiul Independenței 296, 060031 Bucharest, Romania, and <sup>5</sup>Ovidius University Constanta, Faculty of Natural Sciences, Bv. Mamaia 124, Romania

Date submitted: 15 December 2009; Date accepted: 14 April 2010; First published online: 27 July 2010

## SUMMARY

Spatial models are increasingly employed to help understand the distribution of organisms and establish conservation priorities. Classic patch-orientated models may have limited power to accurately predict the organisms' distributions. Pond breeding amphibians are appropriate study organisms because of their complex life cycle, low dispersal and sensitivity to environmental conditions. Here connectivity metrics and niche modelling were used to predict the occurrence of the northern crested newt in a rural landscape from central Romania. Pond-related variables, such as macrophyte cover and the presence of predatory fish, were the most important predictors of newt occurrence, followed by one landscape-related variable (urbanization) and a connectivity metric (nearest neighbouring occupied pond). Most of the landscape and connectivity variables were not adequate predictors, presumably because most of the terrestrial habitats in this traditionally used rural landscape are ecologically optimal for amphibians. Conservation measures for the northern crested newt should promote the preservation of traditional extensive agricultural practices and discourage stocking of ponds with predatory fish.

*Keywords:* amphibian, conservation, habitat, matrix, occurrence, patch, Romania

## INTRODUCTION

The two main approaches to consider spatial patterns of the distribution of organisms in natural systems are landscape ecology (for example Tischendorf & Fahrig 2000) and metapopulation theory (for example Hanski & Gilpin 1991; Hanski 1998). In their classical form, both approaches consider

clearly delimitable patches to predict distribution and spatial dynamic of organisms (Fischer *et al.* 2004). However, patch-based models may be overly simplistic in their prediction of occurrence patterns (Fischer *et al.* 2004; Price *et al.* 2009) and ignoring matrix conditions between patches may lead to serious error (see Marsh & Trenham 2001; Prugh *et al.* 2008; Price *et al.* 2009).

Pond-breeding amphibians represent a challenging group for patch-based analytical approaches of distribution. By their very nature ponds are landscape elements with a particular fit to the metapopulation or 'patch model' approach (Marsh & Trenham 2001). Population size, population turnover and spatial dynamics are often linked to pond quality, primarily through reproductive success (for example Gill 1978; Sjögren-Gulve 1994; Tryjanowski *et al.* 2008). Pond connectivity metrics are often applied in amphibian spatial ecology studies to evaluate the impact of potential source populations and alternative breeding sites on the use of the focal ponds. Four such metrics have been used in 42 European and North American studies (Table 1).

While the distribution and dynamics of some amphibian population systems can be appreciated from the metapopulation perspective (see Gill 1978), elsewhere the landscape effect will mask metapopulation effects (Pope *et al.* 2000). In most of these studies, 'landscape' is represented by the various terrestrial areas (land covers) around the potential breeding pond. Pond-breeding amphibians require terrestrial areas to complete their life history and to replenish their energy reserves by foraging (for example Gibbons 2003). These areas are often partitioned in habitat and non-habitat (or 'matrix'; see Ray *et al.* 2002). The habitat composition of an area, together with the source-sink character of the local population will often be complex and the delimitation of the habitat-population borders is therefore difficult (Thomas & Kunin 1999). The range at which the landscape may affect the status of a local amphibian population is most commonly studied with concentric circles of varying size over which the landscape composition is described. The area does not normally exceed a radius of 4000 m, in line with direct observations on the dispersal range of individuals (Semlitsch 2008). This approach is attractive because it allows the survey of a large number of water bodies (potential breeding habitats)

\*Correspondence: Dr Tibor Hartel Tel/Fax: +40 265 77600 e-mail: asobeka@gmail.com

**Table 1** Survey of pond-based studies ( $n = 42$ ) exploring the distribution of amphibian species with one or more concentric buffers to characterize the surrounding landscape, using wetland area, nearest neighbour distance or metapopulation isolation parameters as covariates.  $X_j$  = pond area ( $A_j$ ), population size ( $N_j$ ) or occupancy data ( $O_j$ ) (see details in Materials and Methods). \*Connectivity index  $C_i = \sum_{j \neq i} \exp(-\alpha d_{ij}) X_j$ . Numbers in the Table represent the number of studies.

Covariate	Europe	North America	Total
Wetland coverage	8	20	28
Wetland density	10	7	17
Nearest pond metric	4	12	16
Connectivity index*	2	1	3

(for example Arntzen & Wallis 1991; Houlihan & Findlay 2003; Ficetola & De Bernardi 2004; Pellet *et al.* 2004; Van Buskirk 2005; see Table 1).

Here we combine the classic metapopulation approach (Hanski & Gilpin 1991) with a niche modelling approach (*sensu* Hutchinson 1957; see also Schmidt & Pellet 2005) to explore the occurrence of the northern crested newt (*Triturus cristatus*) in a rural landscape from Transylvania (Romania). The study area is still largely under traditional management, with small patchy areas of subsistence agriculture, little heavy machinery, low pesticide use and use of natural fertilizers (Hartel *et al.* 2010a). This type of landscape is complex, with high biodiversity in many places, frequently including species that are listed as endangered (Cogălniceanu & Cogălniceanu 2010; see also Tschardt *et al.* 2005 for the characterization of this type of landscape). In these conditions, the habitat versus non-habitat (namely the matrix) character of ecological space may not be obvious for amphibians, owing to the intergradations from habitat to matrix (Hartel *et al.* 2010a).

*Triturus cristatus* is declining in parts of its distributional range, mostly due to pond destruction (see Beebee 1997), and is protected in the European Union (Annex II of the Habitats Directive 92/43/EEC); 850 km<sup>2</sup> of our study area is a Natura 2000 Site of Community Interest (ROSCI0227 Sighișoara-Târnava Mare).

## MATERIALS AND METHODS

### Study area and surveys

The study area covers *c.* 4000 km<sup>2</sup>, and is situated in the middle section of the Târnava Mare River. The dominant land use across this hilly area consists of meadows and pastures (*c.* 40%), deciduous forests (30%) and traditionally managed arable lands (15%) (Hartel *et al.* 2010b). Ponds are mostly man-made, created following the regularization of the Târnava Mare in the 1970s and 1980s and by the damming of tributary streams and springs along the adjacent valleys (Hartel 2008). We studied 96 ponds during 2000–2008 and eliminated data on seven ponds that turned out to be of recent origin (created

<10 years ago). We assumed that these newly-created ponds might give an erroneous picture of the effect of connectivity variables. The ponds were inspected three to four times in one particular year, from late February until May for adults and eggs and until August for larvae, by dip netting and by visual inspection both during the day and at night time. Based on repeated surveys we selected egg searching and night visits with torch use as the most efficient method of species recording for most species (T. Hartel *et al.*, personal observations). The survey methodology and effort was detailed in Hartel *et al.* (2010b).

### Environmental covariates and connectivity measures

We recorded several pond variables, including the percentage cover of emergent vegetation (*Typha* sp. and *Phragmites* sp.) and the occurrence of the predatory fish *Perca fluviatilis*, *Squalius cephalus*, *Silurus glanis*, *Esox lucius*, *Stizosteidon lucioperca*, *Lepomis gibbosus*, *Pseudorasbora parva* and *Clarias gariepinus*. The last three species are invasive; *Clarias gariepinus* has not been reported previously in this area (Hartel *et al.* 2007).

Landscape variables recorded were the percentage cover of meadows and pastures, forests, arable land and urban areas, and the presence/absence of roads with up to 300 cars per hour at night time. These variables were recorded for a radius up to 800 m around each pond. We also noted the distance from each pond to the nearest forest area (in metres) and whether this connection was continuous or interrupted by the presence of presumed barriers, such as arable land, roads and urbanization.

Connectivity was measured as (1) the distance (in metres) to the nearest pond, (2) the occupied state of nearest pond and (3) a species specific connectivity index  $C_i = \sum_{j \neq i} \exp(-\alpha d_{i,j}) O_j$  (modified after Moilanen & Nieminen 2002; see also Pellet *et al.* 2007). Here  $O_j$  is an indicator variable representing the occupancy state of the pond  $j$  (namely  $O_j = 1$  if the  $j$ th pond is occupied and  $O_j = 0$  if not). Moilanen and Nieminen (2002) used the parameter  $A_j^b$  for the area of patch  $j$ , where  $b$  is usually set to 1. Pellet *et al.* (2007) substituted  $A_j^b$  with  $N_j$ , representing the population size in the  $j$ th patch. We used  $O_j$  because it was impossible for us to quantify the population size of the newts and the permanent pond area does not affect the occupancy by newts (Hartel 2008).  $d_{ij}$  is the distance between ponds  $i$  and  $j$ , and  $\alpha$  is the inverse of the mean movement distance. Following Joly *et al.* (2001), we selected a movement distance of 400 m for *T. cristatus*, therefore  $\alpha = 1/400$ .

### Data analysis

We used logistic regression to model the presence of *Triturus cristatus* as a function of the recorded covariates. Prior to the modelling, we examined the data structure for possible non-linearity with the help of additive methods and smoothing functions. We also examined the occurrence of

multicollinearity. We based our model selection procedure on the guidelines of Sauerbrei and Schumacher (1992) and Austin and Tu (2004). The presence of higher order terms and interactions was conditioned on the presence of main effects in the model. From the original data set we drew 1000 non-parametric bootstrap resamples. For each resample we ran a stepwise regression automated model selection procedure. This was a combination of both backward and forward selection, testing at each stage if a variable should be included or excluded from the model. As selection criteria, we chose the Akaike information criterion (AIC), as this is the most common all-purpose model selection tool. AIC is based on the likelihood function, and it seeks the model that best approximates the unknown truth. Information theoretic variable selection differs from that based on significance testing in the sense that non-significant variables can be retained in the model. For each bootstrap iteration, we noted whether a variable was kept in the model, the sign of the regression coefficient and its *p*-value. Data were summarized at 1000 iterations. True predictors were expected to be selected in a high proportion (>50% of iterations), to have a stable sign and to be significant most of the time. After a final model was selected we further examined the effect of removing individual variables. We used receiver operating characteristic curve and the associated area under the curve (AUC) values to assess the predictive powers of the models.

**RESULTS**

Five predictors were identified as possible explanatory factors of the newt presence/absence (Table 2): macrophyte cover, urban areas, predatory fish, road and the nearest pond occupied. The effect of roads was marginally significant and the reduced model (namely without roads) had similar goodness of fit ( $\Delta_{AIC} = 1.228$ ,  $\chi^2 = 3.228$ ,  $p = 0.07$ ) and greater predictive power (AUC = 0.90 with roads and AUC = 0.89 without roads). The removal of roads was, however, beneficial from a modelling perspective because it reduced the stratification of the data and offered more stable and better estimates with lower uncertainty.

No other predictor variables could be removed from the model without compromising its goodness of fit (Table 3). Of the remaining predictors, macrophyte coverage and the nearest occupied pond increased the likelihood of newt presence (Table 4). The effect of macrophyte coverage turned negative after reaching a threshold (*c.* 60%; Fig. 1). We did not find any interaction between the macrophyte cover and predatory fish in determining the pond occupancy by newts. Moreover, no interaction was found between the distance of the closest pond and the occupied state of the closest pond, in determining the occurrence of the newts in our study area. The scenarios suggest that the effect of urban areas on the probability of newt presence was stronger when there were predatory fish in the focal pond and when the neighbouring

**Table 2** Sensitivity analysis based upon non-parametric bootstrapping for the stepwise model selection method incorporating the Akaike information criterion. All values are percentages. Positive values are shown in bold and negative values in italic.

<i>Covariate</i>	<i>Selected</i>	<i>Coefficient sign</i>	<i>Significance</i>
Macrophyte cover	93.3	<b>99.7</b>	88.4
Urban areas	91.8	<i>99.8</i>	82.7
Predatory fish	86.8	<i>99.5</i>	83.1
Macrophyte cover <sup>2</sup>	67.4	<i>93.0</i>	69.4
Road	57.8	<i>94.1</i>	57.6
Nearest pond occupied	53.2	<b>91.9</b>	55.8
Arable land cover	49.9	<i>85.4</i>	67.1
Pond-forest connection	43.5	<i>81.6</i>	48.7
Distance from forest	38.2	<i>57.3</i>	37.7
Grassland/pasture cover	35.0	<i>84.3</i>	43.7
Nearest pond distance	30.8	<i>80.5</i>	40.3
Forest cover	28.5	<i>54.0</i>	39.7
Connectivity index	19.0	<i>74.2</i>	32.6

**Table 3** Influence of variable removal from the whole model on the information content (AIC) and predictive power (AUC). Likelihood ratio test (LRT) against the removal is also provided. When the macrophyte coverage was removed automatically, the second order term was also removed.

<i>Variable</i>	<i>df</i>	<i>AUC</i>	<i>AIC</i>	<i>LRT</i> $\chi^2$	<i>p</i>
No removal	1	0.891	80.28		
Macrophyte cover	1	0.827	88.01	9.72	0.001
Predatory fish	1	0.856	89.49	11.20	0.0008
Urban area cover	1	0.871	87.50	9.21	0.002
Nearest pond occupied	1	0.872	83.09	4.80	0.02
Macrophyte cover <sup>2</sup>	1	0.875	83.44	5.16	0.02


**Table 4** Contribution of five selected environmental factors to the model that explores the occurrence of the northern crested newt (*Triturus cristatus*) in 89 ponds in central Romania.

<i>Variable</i>	<i>B</i>	<i>SE</i>	<i>Z value</i>	<i>Pr</i> (> z )
Intercept	-1.55	0.82	-1.88	0.060
Macrophyte cover	0.10	0.03	2.84	0.004
Predatory fish	-2.32	0.75	-3.08	0.002
Urban area cover	-6.98	2.95	-2.36	0.018
Nearest pond occupied	1.35	0.62	2.16	0.030
Macrophyte cover <sup>2</sup>	-0.0003	0.0003	-2.211	0.027

pond was unoccupied than in the reverse situation (Fig. 1).

**DISCUSSION**

Both metapopulation and niche models contributed to understanding the distribution of northern crested newt in the rural landscape of central Romania. We found that pond-related variables (predatory fish and macrophyte cover) were the most important determinants of *T. cristatus* distribution in the study area, followed by one landscape-related variable



**Figure 1** Probability of the presence of the northern crested newt (*Triturus cristatus*) as a function of pond macrophyte coverage and the degree of urbanization within an 800 m radius of the pond, in ponds with (*b, d*) and without fish (*a, c*), and the nearest unoccupied pond (*a, b*) versus those occupied by *T. cristatus* (*c, d*) in 89 ponds in central Romania.

(the urban area cover), one spatial metric (the nearest occupied pond) and the quadratic effect of macrophyte cover. Six out of seven landscape covariates and two out of three connectivity variables were not found to be important for *T. cristatus*. Because human settlements and ponds, through their rapid change, represent the two most significant impacts in this rural area over the last decades, these results were not unexpected.

Increasing private ownership of ponds has resulted in increased pond stocking with both greater numbers of fish species and especially more predatory fish species. *Triturus cristatus* is sensitive to fish presence (for example Joly *et al.* 2001; Van Buskirk 2005; Skei *et al.* 2006; Gustafson *et al.* 2009; Kloskowski 2010). Features that expose *T. cristatus* to fish predation include a long aquatic period both as adults (breeding and feeding) and larvae (Malmgren 2001) and pelagic larvae that are easily spotted by predators, with no behavioural avoidance in the presence of aquatic predators. Also, the ability of northern crested newt larvae to respond to predators by accelerating larval development is very limited (Schmidt & Van Buskirk 2001). This inability to reduce predation risk by morphological plasticity may further expose *T. cristatus* larvae to fish predation.

The emergent vegetation cover influenced the probability of newts occurring up to c. 60% coverage. A similar threshold of macrophyte cover on pond occupancy was found by Oldham *et al.* (2000) for *T. cristatus* and on newt abundance in general by Joly *et al.* (2001). The rationale for the effect is probably the lack of available space for newt breeding and foraging activity (Oldham *et al.* 2000). Similarly, Gustafson *et al.* (2009) found

that ponds in late succession stages had become inhospitable for *T. cristatus* and suggested management and restoration activities to keep ponds open. Late succession stages also negatively influenced pond use by *Rana lessonae* and *R. dalmatina* (Sjögren Gulve 1994; Hartel *et al.* 2009). We found no significant interaction between the macrophyte coverage and presence of predatory fish, suggesting that emergent vegetation will not buffer the effect of predators. This may be because small predatory fish (such as *P. parva*, *L. gibbosus*, *P. fluviatilis*) use the macrophyte as breeding area and fish fry may use it as a refuge area.

The distance to the next nearest occupied pond was the only spatial predictor of newt presence, confirming that simple linear connectivity measures can be useful for predicting animal distribution and may even outperform more complex metrics (Prugh 2009). The independent effect of newt presence in the next closest pond from the distance to the closest pond suggests that isolation by distance is not a problem in these landscapes (but see Halley *et al.* 1996). The connectivity index  $C_i$  had no effect on newt presence in the focal ponds, possibly because of the dichotomous character of  $O_j$ , as opposed to a variable that denotes area and/or population size of the potential sources (see Moilanen & Nieminen 2002; Pellet *et al.* 2007). This may lead to the loss of important information, because both population size and distance contribute to the persistence of a newt metapopulation network. Halley *et al.* (1996) found that the likelihood of the persistence of newt ponds increased if ponds supported >40 females, or were <0.5 km from a typical source

pond. The likelihood of persistence of small newt ponds (<10 females) depended on the existence of a source pond < 0.75 km away, whereas for large newt ponds a source pond could be <1.5 km away (Halley *et al.* 1996).

As expected, the landscape composition had little effect on newt occurrence, suggesting that unsuitable terrestrial habitats (other than urban area cover) are not currently a limiting factor for the newt (see also Hartel *et al.* 2010a for other amphibians). Terrestrial habitat degradation may result in increased pond to pond isolation. In our study area, newly-created permanent ponds were colonized by *T. cristatus* within one or two years (Hartel & Öllerer 2009). This may reflect the unaffected nature of the rural landscape because, in intensively-managed landscapes, the colonization capability of *T. cristatus* was lower than that of most other amphibian species (for example Laan & Verboom 1990; Beebee 1997).

## CONCLUSIONS

We have shown that the occurrence of the northern crested newt can be understood using the combination of metapopulation and niche modelling approaches; occurrence in a traditionally managed rural area is primarily related to pond variables. Avoidance of predatory fish introduction and management of aquatic vegetation density should be priority activities for northern crested newt conservation in this area. Fish removal from stocked ponds may also be an appropriate management implication, and such an activity may be encouraged especially in the area covered by the Natura 2000 regulations. No unequivocal results were found regarding the importance of terrestrial habitats around ponds, other than a negative effect of urbanization. Hence, a dense metapopulation network for *T. cristatus* would be composed of many permanent ponds distributed over a non-urban extensively-managed landscape. In order to better understand the connectivity between ponds in relation to pond management and land management in a cultural landscape, more studies are needed on the neighbourhood size and dispersal distance of the northern crested newt and other amphibians. We also suggest the fragment-based spatial models as employed in this study be complemented with alternative models that consider gradients between landscape elements (Fischer *et al.* 2004), because these may be especially useful in complex traditionally-managed landscapes such as found in Romania and other parts of Eastern Europe.

## ACKNOWLEDGEMENTS

This study was conducted with the financial support from the Declining Amphibian Populations Task Force and the Swedish Biodiversity Centre. We thank Dr. Jerome Pellet for his advice in calculating the connectivity metric and three anonymous reviewers for their constructive comments on the manuscript.

## References

- Arntzen, J.W. & Wallis, G.P. (1991) Restricted gene flow in a moving hybrid zone of the newts *Triturus cristatus* and *T. marmoratus* in Western France. *Evolution* **45**: 805–826.
- Austin, P. & Tu, J. (2004) Bootstrap methods for developing predictive models. *The American Statistician* **58**: 131–137.
- Beebee, T.J.C. (1997) Changes in dewpond numbers and amphibian diversity over 20 years on chalk downland in Sussex, England. *Biological Conservation* **81**: 215–219.
- Cogălniceanu, D. & Cogălniceanu, G.C. (2010) An enlarged European Union challenges priority settings in conservation. *Biodiversity and Conservation* **19**: 1471–1483.
- Ficetola, G.F. & De Bernardi, F. (2004) Amphibians in a human dominated landscape: the community structure is related to habitat features and isolation. *Biological Conservation* **119**: 219–230.
- Fischer, J., Lindenmayer, D.B. & Fazey, I. (2004) Appreciating ecological complexity: habitat contours as a conceptual landscape model. *Conservation Biology* **18**: 1245–1253.
- Gibbons, J.W. (2003) Terrestrial habitat: a vital component for herpetofauna of isolated wetlands. *Wetlands* **23**: 630–635.
- Gill, D.E. (1978) The metapopulation ecology of the red spotted newt, *Notophtalmus viridescens* Rafinesque. *Ecological Monographs* **48**: 145–166.
- Gustafson, D.H., Andersen, A.S.L., Mikusinski, G. & Malmgren, J.C. (2009) Pond quality determinants of occurrence patterns of northern crested newts (*Triturus cristatus*). *Journal of Herpetology* **43**: 300–310.
- Halley, J.M., Oldham, R. & Arntzen, J.W. (1996) Predicting the persistence of amphibian populations with the help of a spatial model. *Journal of Applied Ecology* **33**: 455–470.
- Hanski, I. (1998) Metapopulation dynamics. *Nature* **396**: 41–49.
- Hanski, I. & Gilpin, M. (1991) Metapopulation dynamics: brief history and conceptual domain. *Biological Journal of the Linnean Society* **42**: 3–16.
- Hartel, T. (2008) Ecological and faunistic studies on amphibians in the Tarnava Mare basin. Ph.D. dissertation, Ovidius University, Constanta, Romania.
- Hartel, T. & Öllerer, K. (2009) Local turnover and factors influencing the persistence of amphibians in permanent ponds from the Saxon landscapes of Transylvania. *North-Western Journal of Zoology* **5**: 40–52.
- Hartel, T., Nemes, Sz., Cogălniceanu, D., Öllerer, K., Schweiger, O., Moga, C.I. & Demeter, L. (2007) The effect of fish and habitat complexity on amphibians. *Hydrobiologia* **583**: 173–182.
- Hartel, T., Nemes, Sz., Cogălniceanu, D., Öllerer, K., Moga, C.I., Lesbarreres, D. & Demeter, L. (2009) Pond and landscape determinants of *Rana dalmatina* population sizes in a Romanian rural landscape. *Acta Oecologica* **35**: 53–59.
- Hartel, T., Öllerer, K., Cogălniceanu, D., Nemes, Sz., Moga, C.I. & Demeter, L. (2010a) Pond-based survey of amphibians in a Saxon cultural landscape from Transylvania (Romania). *Italian Journal of Zoology* **77**: 61–70.
- Hartel, T., Schweiger, O., Öllerer, K., Cogălniceanu, D. & Arntzen, J.W. (2010b) Amphibian distribution in a traditionally managed rural landscape of Eastern Europe: probing the effect of landscape composition. *Biological Conservation* **143**: 1118–1124.
- Houlahan, J.E. & Findlay, C.S. (2003) The effects of adjacent land use on wetland amphibian species richness and community composition. *Canadian Journal of Fisheries and Aquatic Sciences* **60**: 1078–1094.

- Hutchinson, G. E. (1957) Concluding remarks. Population studies: animal ecology and demography. *Cold Springs Harbor Symposium on Quantitative Biology* 22: 415–427.
- Joly, P., Miaud, C., Lehmann, A. & Grolet, O. (2001) Habitat matrix effect on pond occupancy in newts. *Conservation Biology* 15: 239–248.
- Kloskowski, J. (2010) Fish farms as amphibian habitats: factors affecting amphibian species richness and community structure at carp ponds in Poland. *Environmental Conservation* 37 DOI:10.1017/S0376892910000494
- Laan, R. & Verboom, B. (1990) Effects of pool size and isolation on amphibian communities. *Biological Conservation* 54: 251–262.
- Malmgren, J.C. (2001) Evolutionary ecology of newts. Ph.D. dissertation. Studies in Biology 1, Örebro University, Örebro, Sweden.
- Marsh, D.M. & Trenham, P.C. (2001) Metapopulation dynamics and amphibian conservation. *Conservation Biology* 15: 40–49.
- Moilanen, A. & Nieminen, M. (2002) Simple connectivity measures in spatial ecology. *Ecology* 83: 1131–1145.
- Oldham, R.S., Keeble, J., Swan, M.J.S. & Jeffcote, M. (2000) Evaluating the suitability of habitat for the northern crested newt (*Triturus cristatus*). *Herpetological Journal* 10: 143–155.
- Pellet, J., Fleishman, E., Dobkin, D.S., Gander, A. & Murphy, D.D. (2007) An empirical evaluation of the area and isolation paradigm of metapopulation dynamics. *Biological Conservation* 136: 483–495.
- Pellet, J., Guisan, A. & Perrin, N. (2004) A concentric analysis of the impact of urbanization on the threatened European tree frog in an agricultural landscape. *Conservation Biology* 18: 1599–1606.
- Pope, S.E., Fahrig, L. & Merriam, N.G. (2000) Landscape complementation and metapopulation effects on leopard frog populations. *Ecology* 81: 2498–2508.
- Price, B., McAlpine, C.A., Kutt, A.S., Phinn, S.R., Pullar, D.V. & Ludwig, J.A. (2009) Continuum or discrete patch landscape models for savanna birds? Towards a pluralistic approach. *Oikos* 32: 745–756.
- Prugh, L.R. (2009) An evaluation of patch connectivity measures. *Ecological Applications* 19: 1300–1310.
- Prugh, L.R., Hodges, K.E., Sinclair, A.R.E. & Brashares, J.J. (2008) Effect of area and isolation on fragmented animal populations. *Proceedings of the National Academy of Sciences* 105: 20771–20775.
- Ray, N., Lehmann, A. & Joly, P. (2002) Modeling spatial distribution of amphibian populations: a GIS approach based on habitat matrix permeability. *Biodiversity and Conservation* 11: 2143–2165.
- Sauerbrei, W. & Schumacher, M. A. (1992) A bootstrap resampling procedure for model building: application to the Cox regression model. *Statistics in Medicine* 11: 2093–2109.
- Schmidt, B.R. & Pellet, J. (2005) Relative importance of population processes and habitat characteristics in determining site occupancy of two anurans. *Journal of Wildlife Management* 69: 884–893.
- Schmidt, B.R. & Van Buskirk, J. (2001) Predator induced phenotypic plasticity in larvae of crested newt, *Triturus cristatus*. *Sonderheft* 4: 179–191.
- Semlitsch, R.D. (2008) Differentiating migration and dispersal processes for pond-breeding amphibians. *Journal of Wildlife Management* 72: 260–267.
- Sjögren-Gulve, P. (1994) Distribution and extinction patterns within a northern metapopulation of the pool frog, *Rana lessonae*. *Ecology* 75: 357–367.
- Skei, J.K., Dolmen, D.L., Rønning, L. & Ringsby, T.R. (2006) Habitat use during the aquatic phase of the newts *Triturus vulgaris* (L.) and *T. cristatus* (Laurenti) in central Norway: proposition for a conservation and monitoring area. *Amphibia-Reptilia* 27: 309–327.
- Thomas, C.D. & Kunin, W.E. (1999) The spatial structure of populations. *Journal of Animal Ecology* 68: 647–657.
- Tischendorf, L. & Fahrig, L. (2000) On the usage and measurement of landscape connectivity. *Oikos* 90: 7–19.
- Tryjanowski, P., Sparks, T., Rybacki, M. & Berger, L. (2006) Is body size of the water frog *Rana esculenta* complex responding to climate change? *Naturwissenschaften* 93: 110–113.
- Tscharntke, T., Klein, A.M., Steffan-Dewenter, I. & Thies, C. (2005) Landscape perspectives on agricultural intensification and biodiversity: ecosystem service management. *Ecology Letters* 8: 857–874.
- Van Buskirk, J. (2005) Local and landscape influence on amphibian occurrence and abundance. *Ecology* 86: 1936–1947.
- Zanini, F., Pellet, J. & Schmidt, B.R. (2009) The transferability of distribution models across regions: an amphibian case study. *Diversity and Distributions* 15: 469–480.